If you have dental implants and a doctor orders an MRI, it is completely reasonable to pause and ask whether those two things can safely coexist. Patients ask this often, usually right after hearing the words "magnetic resonance imaging." The concern makes sense. An MRI uses a powerful magnetic field, and most people have at least a vague sense that metal and magnets are not always a friendly combination.

The short answer is yes, in most cases you can get an MRI with dental implants. For the vast majority of patients, dental implants do not prevent MRI scanning and do not create a safety hazard. That said, the full answer deserves more nuance than a simple yes or no. The implant material matters. The body area being scanned matters. The age and design of the restoration can matter. Even when an MRI is safe, it can still produce image distortion in some situations.

This is one of those medical questions where precision helps. "Dental implants" can refer to the titanium post in the jawbone, the abutment that connects the post to the visible tooth, the crown itself, or in some cases additional components such as bars, magnets, or removable overdenture attachments. Not all metal in the mouth behaves the same way in the MRI environment.

Understanding what is actually in your mouth, and what your imaging team needs to know, can save time and prevent unnecessary worry.

Why dental implants usually are not a problem

Most modern dental implants are made from titanium or titanium alloys. Titanium is widely used in medicine for good reasons. It is strong, corrosion resistant, biocompatible, and importantly for this discussion, it is not ferromagnetic in the way iron, nickel, or certain steels can be. In practical terms, that means it is not strongly pulled by the MRI magnet.

That is the key point. The nightmare scenario many patients imagine is that the implant could be ripped out or shift inside the jaw. With a standard, well integrated titanium implant, that is not how this works. Once the implant has fused with the surrounding bone through osseointegration, it is mechanically stable. The magnetic field of the MRI does not suddenly overpower that bond and yank the implant loose.

In clinical practice, people with dental implants undergo MRI scans every day. Brain MRIs, spine MRIs, knee MRIs, abdominal MRIs, pelvic MRIs, these are routine. If a patient has a healed titanium implant and no other complicating oral hardware, the implant itself is generally not the reason to cancel or avoid the scan.

That reassurance is important, but it should not slide into carelessness. Safe is not identical to irrelevant. Dental implants can still affect MRI images, especially when the scan is being performed near the mouth, face, or jaw.

Safety versus image quality, two different questions

A lot of confusion comes from mixing up two separate issues. One is whether the MRI is safe to perform. The other is whether the implant will interfere with the images.

From a safety standpoint, standard titanium dental implants are usually compatible with MRI. From an image quality standpoint, they can create artifact, which is a distortion or shadowing effect in the scan. The metal can disrupt the local magnetic field enough to blur or obscure nearby structures.

This matters most when the area of interest is close to the implant. If someone needs an MRI of the ankle and has a dental implant, image distortion from that implant is not likely to matter. If someone needs an MRI of the temporomandibular joint, tongue, salivary glands, sinuses, inner ear region, or brainstem area, artifact from oral metal may be more relevant.

Radiologists and MRI technologists deal with this routinely. Sometimes the artifact is minor and acceptable. Sometimes they can modify the imaging sequence to reduce it. Sometimes a different imaging test, such as CT, cone beam CT, ultrasound, or another MRI protocol, may provide better information depending on the clinical question.

In other words, the presence of dental implants usually does not stop the scan, but it may influence how the scan is planned and how useful the final images are.

What most implants are made of

When people say "metal implant," they often imagine a single category. In reality, dental restorations can include several materials.

The implant fixture, the part placed in the jawbone, is most often titanium. Some implants are made from zirconia, which is a ceramic, not a metal. Zirconia implants are not magnetic and generally create less artifact than many metals, although imaging behavior can still vary depending on the system and surrounding materials.

The visible restoration on top of the implant may be porcelain, zirconia, metal ceramic, or another material. The abutment may be titanium, zirconia, or a combination. If the patient wears an implant supported overdenture, the attachment mechanism may include metal housings or in some designs small magnetic components. Those details can matter more than patients realize.

This is one reason imaging centers ask broad questions about "implants," "bridges," "dentures," and "oral appliances." They are not trying to be difficult. They are trying to identify which item is fixed, which is removable, and which could affect either safety or image quality.

Cases that deserve a closer look

Although the typical titanium dental implant is not a contraindication to MRI, there are situations where the answer is less automatic.

Older dental work can be Dental Implants harder to identify, especially if the patient does not have records and the implant was placed many years ago. Some older restorations may include alloys or design features that are not immediately obvious from memory alone. That does not mean they are dangerous, only that the MRI team may want more information before proceeding.

Removable implant overdentures sometimes need special attention. The implants in the bone may stay in place, but the denture itself, if removable, may need to come out before the scan. This is true for many oral appliances, not because they are all unsafe, but because they can create artifact, heat, discomfort, or movement. Any appliance with magnets deserves explicit review.

Patients who have had head and neck cancer treatment may have a more complex reconstruction history, such as plates, screws, grafts, obturators, or surgical hardware. In those cases, the phrase "I have dental implants" may capture only part of the picture. The MRI facility should know about all prior oral and facial procedures.

There is also the issue of field strength. Many routine clinical scanners operate at 1.5 Tesla or 3 Tesla. In general, common dental implants are managed safely at these standard field strengths, but artifact may be more pronounced at higher strengths. Protocol decisions become more important when the anatomy of interest sits close to the implant.

What patients often feel during an MRI

One practical worry patients mention is whether they will feel the implant heating up or vibrating. Significant heating of a standard, fixed dental implant is not what is typically seen in routine MRI settings, but comfort can vary and unusual sensations should always be reported. Some patients notice a strange awareness of their dental work during scanning, though often that anxiety fades once the test begins and nothing happens.

The louder concern, usually, is not the implant itself but the experience of MRI. It is noisy. It can feel confined. It requires staying still. If the scan is of the head or neck, the coil around the face can make people feel especially enclosed. Those factors tend to bother patients far more than the presence of dental implants.

When I have seen this discussed clinically, the patients who struggle most are rarely those at risk from the implant. They are the ones who were not told what to expect, or who arrived assuming any metal in the body must be dangerous. Clear communication solves much of that fear.

Why your radiology center asks so many questions

MRI screening forms can feel repetitive. You may be asked about implants, surgeries, crowns, braces, retainers, dentures, hearing devices, piercings, and old injuries involving metal fragments. The staff may ask those same questions again in person, even after you filled out the form online.

That repetition is deliberate. MRI safety depends on details, and details are often lost when people use casual labels. One patient may call a removable partial denture an implant. Another may say "crown" when they actually have a post and core, bridge, and implant under it. Another may forget a magnetic denture attachment entirely because it has been in place for years and feels like part of them.

If you know where and when your dental implants were placed, bring that information. If you have an implant card or records from your dentist or oral surgeon, even better. Many modern implant systems come with documentation that identifies the manufacturer and material. The MRI team may not need it, but if there is any question, it can speed things up.

MRI of the brain, face, and jaw, where artifact matters most

When the target anatomy lies above the shoulders, the discussion becomes more specific. Dental implants can create streaking, signal voids, and local distortion, especially on some sequences. The impact varies by implant composition, number of implants, proximity to the anatomy being evaluated, and the technical choices made by the imaging team.

A single posterior implant in the lower jaw may have negligible effect on a brain MRI focused on structures high in the skull. Multiple implants, extensive fixed prosthetics, or metal rich restorations across both arches can cause broader artifact that spills into adjacent regions. This does not always ruin the study, but it can obscure pathology in nearby tissues.

For patients being evaluated for oral cancer, salivary gland disease, jaw lesions, or trigeminal symptoms, that distinction matters. A scan may still be worth doing, yet the ordering physician and radiologist may need to balance MRI's strengths against its limitations. In some cases, CT shows bone better. In others, MRI is still preferred for soft tissue detail despite the artifact. The right test depends on the question being asked.

That is why a blanket internet answer can be misleading. "Dental implants are MRI safe" is broadly true. "Dental implants never affect MRI" is not.

Braces, crowns, fillings, and implants are not the same thing

Patients often group all dental hardware together, but MRI compatibility is not a single category. Orthodontic braces are a different issue from dental implants. Braces are more likely to interfere significantly with imaging of the head and face because they sit on the tooth surfaces across a wide area. Fillings and crowns vary by material and usually create less concern, though they can still produce artifact.

Dental implants tend to be deeply anchored in bone and usually made of MRI compatible material, which is reassuring from a safety standpoint. The restoration on top of the implant may add some artifact, but the underlying implant itself is generally not the dramatic problem people picture.

This distinction matters because some patients hear that "metal in the mouth affects MRI" and assume that means all oral metal is equally dangerous. It does not. The real conversation is about type, location, magnetism, and imaging effect.

What you should tell the MRI team

The safest approach is simple: disclose everything you know about your dental work, even if you suspect it is ordinary. Mention fixed implants, removable implant supported dentures, clips, bars, magnetic attachments, and any oral appliance you wear regularly.

A good practical checklist looks like this:

1. Tell the staff you have dental implants and where they are located.
2. Mention whether anything in your mouth is removable.
3. Bring implant documentation if you have it, especially for older or uncommon systems.
4. Report any prior head, neck, or jaw surgery beyond routine implant placement.
5. Ask whether the scan area is close enough to the implants that artifact could affect results.

This is not about making the process harder. It is about giving the technologist and radiologist the information they need before the scan starts, not halfway through when an image comes back distorted.

If you are missing records, do not panic

A common real world problem is incomplete dental history. People move, dentists retire, offices close, records disappear, and ten years later an imaging center asks for details nobody remembers. That can be frustrating, but it does not automatically stop the MRI.

Often the facility can proceed based on the most likely scenario, particularly if the implants are modern and fixed. If there is genuine uncertainty, they may contact your dentist, review prior imaging, or consult the radiologist and referring physician. In some cases, a plain x ray or CT already on file helps clarify what type of hardware is present.

The key is not to guess confidently if you are unsure. Saying "I think they are titanium, but I do not know for certain" is far more useful than saying "there is nothing metal in my mouth" when you do in fact have implants. Good MRI screening depends more on honesty than on perfect memory.

The rare situations when MRI may need to be delayed or adjusted

It is uncommon for standard Dental Implants alone to block MRI, but delays can happen when there is uncertainty about additional hardware or when removable components have not been accounted for. A patient may arrive wearing an implant retained overdenture and not realize it should be removed. Another may have a facial prosthesis with small magnets. Another may have had reconstructive surgery and forgotten to mention plates or clips.

Sometimes the scan itself is possible, but the protocol needs adjustment. Metal artifact reduction techniques may help. Positioning changes may help. A different sequence may help. Occasionally the radiologist may advise a different imaging modality if the expected artifact is likely to make the MRI less informative.

These are workflow issues, not signs that something has gone wrong. MRI is a highly adaptable tool, but adaptation works best when the imaging team knows what they are dealing with.

What about newly placed implants?

Patients sometimes worry more if the implant was placed recently and has not fully healed. From a practical standpoint, the material properties of the implant do not change just because it is new. The bigger concern in the early postoperative period is usually comfort, not magnetic attraction. Lying still for a long scan may be unpleasant if the surgical site is tender. If there is any question, the oral surgeon or restorative dentist can advise based on the timing of placement and your healing progress.

It is also worth distinguishing an implant from temporary healing components or provisional prostheses. Those parts may differ in material and removability. If you are in the middle of implant treatment rather than years out from it, give the imaging center that context.

A useful way to think about the risk

For most patients, a dental implant is more like a hip replacement or a fixation screw than a loose piece of metal jewelry. It is a medical device chosen in part because it behaves predictably in the body. That does not mean every implant restoration is identical, but it explains why MRI and dental implants are so commonly compatible.

The more meaningful question is often not "Can I have the MRI?" But "Will my dental work interfere with the answer my doctor needs from the MRI?" That is a subtler and better question. It shifts the conversation from fear to planning.

If the scan is for the knee, your implant is almost certainly an afterthought. If the scan is for the jaw joint or oral cavity, your implant may become part of the technical discussion. Same patient, same implant, different clinical context.

The bottom line for patients with dental implants

Most people with dental implants can safely undergo MRI. Modern implants are usually made from titanium or similarly compatible materials, and they are firmly integrated into bone. They are not typically pulled by the magnet or made dangerous by the scan.

Where dental implants become more relevant is image quality, especially for MRI studies of the head, face, mouth, and nearby structures. Artifact can limit what the radiologist sees, and that possibility should be considered before the scan, not discovered by surprise afterward.

If you have Dental Implants and need an MRI, tell the imaging center in advance, share whatever records you have, and mention any removable oral components or prior facial surgery. That small step usually makes the process smooth. In most cases, the answer is reassuring: yes, you can get the MRI, and with the right information, your care team can decide how to get the best images possible.

Oaks Dental
5000 Parkway Calabasas Suite 308
Calabasas, CA 91302, United States
+1 818-431-2000

FAQ About Dental Implants

What is the downside of a dental implant?

Implants require surgery and a healing period that can last several months. Possible complications include infection, nerve or sinus injury, damage to nearby teeth and failure to bond with bone. Your dentist should assess your health, bone support and alternatives before recommending treatment.

What do dental implants typically cost?

There is no single price for every implant case. The number of teeth replaced, implant components, final restoration and any grafting or extractions affect the total. Request a written estimate covering every stage and confirm insurance benefits and out-of-pocket costs with the practice.

How long do tooth implants last?

An implant can provide long-term support and may last a lifetime with good oral hygiene and regular dental care, but this is not guaranteed. The crown or other restoration may need replacement sooner. Smoking, gum disease and some health conditions can reduce implant success.